

Multi-phase separation in mitochondrial nucleoids and eukaryotic nuclei

Qi Long^{1,2}, Yanshuang Zhou^{1,2}, Jingyi Guo^{1,2}, Hao Wu^{1,2}, Xingguo Liu^{1,2,3}✉

¹ CAS Key Laboratory of Regenerative Biology, Joint School of Life Sciences, The Sixth Affiliated Hospital of Guangzhou Medical University, Qingyuan People's Hospital, Guangzhou Medical University; Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China

² Guangdong Provincial Key Laboratory of Stem Cell and Regenerative Medicine, China-New Zealand Joint Laboratory on Biomedicine and Health, CUHK-GIBH Joint Research Laboratory on Stem Cells and Regenerative Medicine, Institute for Stem Cell and Regeneration, Institute for Stem Cell and Regeneration, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China

³ Centre for Regenerative Medicine and Health, Hong Kong Institute of Science & Innovation, Chinese Academy of Sciences, Hong Kong SAR, China

Received: 11 August 2022 / Accepted: 10 February 2023

Abstract In mammalian cells, besides nuclei, mitochondria are the only semi-autonomous organelles possessing own DNA organized in the form of nucleoids. While eukaryotic nuclear DNA compaction, chromatin compartmentalization and transcription are regulated by phase separation, our recent work proposed a model of mitochondrial nucleoid self-assembly and transcriptional regulation by multi-phase separation. Herein, we summarized the phase separation both in the nucleus and mitochondrial nucleoids, and did a comparison of the organization and activity regulating, which would provide new insight into the understanding of both architecture and genetics of nucleus and mitochondrial nucleoids.

INTRODUCTION

The compartmentation in cells is critical for carrying multiple functions (Boeynaems *et al.* 2018). It is well known that the inner membrane system segregates space in cells into organelles, such as the endoplasmic reticulum, Golgi apparatus, lysosomes, endosomes and so on. These organelles concentrate special factors to generate small reaction centers, which provide spatiotemporal control over cellular materials, metabolic and signals for cells in carrying multiple functions and pathways. Despite the inner membrane system, phase separation could also concentrate biomolecule independence of the membrane systems, which greatly expands the compartmentation in cells

and reduces the threshold for generating small reaction centers (Banani *et al.* 2016, 2017; Riback *et al.* 2020).

Phase separation serves as a biomolecular condenser, which concentrates biomolecules to form droplet-like or gel-like structures (Banani *et al.* 2017; Shin and Brangwynne 2017). It is a kind of self-assembly driven by the intrinsic property of the biomolecules, such as weak interaction beneath intrinsically disordered regions (IDR) or multivalent effect between molecule interaction (Alberti *et al.* 2019). Not only proteins, DNAs, RNAs and even small molecules, such as dNTPs and drugs could be concentrated by co-phase separation (Klein *et al.* 2020; Wang *et al.* 2018), which provides the substrates, enzymes or cofactors for the efficient reaction in the small reaction centers generated by phase separation.

Phase separation provides the driving force to concentrate biomolecules into membraneless organelles, such as stress granules, cell junctions, signal

Qi Long and Yanshuang Zhou contributed equally to this work.

✉ Correspondence: liu_xingguo@gibh.ac.cn (X. Liu)

transduction, transcription complex and so on (Banani *et al.* 2016). It is also reported to take part in the condensation of chromatin, enhancers, nuclear speckles, heterochromatin and even the architecture of nuclei, such as the generation of the nucleolus (Sabari *et al.* 2020). Nuclei are the largest organelle in cells and play critical important and multiple roles in DNA storage, replication and transcription, ribosome assembling, *etc.* The compartmentation by phase separation is critical for the nuclei in carrying multiple roles.

Mitochondria are the only semi-autonomous organelle in mammalian cells, owing a circle DNA, which plays critical roles in energy production, metabolism, apoptosis, and so on. Mitochondria DNA is organized into nucleoids by TFAM, one of the most abundant proteins in mitochondria. The crista plays a critical role in the compartmentation of mitochondria for metabolism and energy production (Cogliati *et al.* 2013; Frey and Mannella 2000). However, there are rare reports about the compartmentation in mitochondrial nucleoids, which share many commons with the nuclei, as well as some unique features. Recently, we reported that phase separation drives the self-assembly of the nucleoid, and regulates mitochondrial transcription, which provides a new model for mitochondria compartmentation (Long *et al.* 2021).

Herein, we summarized the compartmentation of nuclei and mitochondria in architecture assembling and discussed the phase separation from nuclei to mitochondrial nucleoids.

MAIN

Phase separation drives the chromatin condensation in nuclear architecture assembly

Nuclei are membrane organelles, which are responsible for chromatin-DNA storage, replication and transcription, *etc.* The chromatin-DNA is compacted by histones and condensates gradually from nucleosomes, 10 nm chromatin fiber, 30 nm chromatin fiber, euchromatin and heterochromatin to chromosomes and stored in the nuclei (Misteli 2020; Song *et al.* 2014). The chromatin-containing enhancers, especially super-enhancers could be gathered together with activators in a phase separation manner, which promotes transcription activity (Sabari *et al.* 2018). What's more, the chromatin fiber could be locked by CTCF loops and clustered into droplets similarly driven by phase separation (Hansen *et al.* 2019). Many nuclear speckles acting as small reaction centers are generated by

chromatin fiber, RNAs and so on, which are also reported to be driven by phase separation (Bi *et al.* 2019; Guo *et al.* 2019). The m6A-modification on RNA is reported to be critical for generating transcription activation speckles by phase separation (Cheng *et al.* 2021; Lee *et al.* 2021). The chromatin fiber could be further condensed by linker histone H1 and HP1a which also drives the special heterochromatin foci formation in a phase separation manner (Larson *et al.* 2017; Strom *et al.* 2017; Wang *et al.* 2020). The phase separation induces the aggregation of nucleosomes or chromatin fibers. What's more, the condensation by phase separation in heterochromatin prevents the invasion of transcription factors, maintaining the inactivation status (Larson and Narlikar 2018; Sabari *et al.* 2020). In contrast, the euchromatin is enriched with transcription activation factors, RNA polymerase, RNAs and so on, but not H1 and HP1a, promoting gene activation and mRNA transcription.

The phase separation also provides a mechanical force for the architecture assembly and maintenance in cell nuclei. The chromatin is reported to be a gel-like structure, rather than liquid as the previous report (Strickfaden *et al.* 2020). The gel-like structure is also a feature of phase separation, which shows properties between solids and liquids. The gel-like structure of chromatin should also provide important mechanical support for the nucleus. The dynamic of the phase separation in chromatin may also regulate the mechanics of the nuclei, resulting in cell migration and invasion (Nava *et al.* 2020).

The nucleolus is the largest sub-organelle in the nuclei, which could be easily visualized under a normal microscope even in a bright field. It is a special reaction center without membrane segregation, which is responsible for rRNA transcription and ribosome assembly. It is reported to be a layered structure, starting from an inner core from rRNA transcription and proceeding toward the periphery (Brangwynne *et al.* 2011; Lafontaine *et al.* 2021). To keep the reaction smoothly and steadily, a multi-layered droplet-like liquid-liquid phase separation was found to drive the nucleolus formation (Feric *et al.* 2016; Lafontaine *et al.* 2021).

Phase separation drives the assembly of the mitochondrial nucleoid

Unlike nuclei, the nucleoid is a membraneless organelle in mitochondria, lacking membranes to segregate it from the mitochondrial matrix. Mitochondria are thought to be derived from ancient bacteria (Roger *et al.* 2017), which possess a simple circle DNA,

compacted by TFAM proteins into a nucleoid structure, similar to the nucleoid in bacteria (Kukat *et al.* 2015). Although the nucleoid is much more relaxed than chromatin, it shares many commons with nucleosomes. The high abundance of TFAM coated mtDNA in a ratio of 1:1000 to 1:2000 as reported (Ngo *et al.* 2014; Shen and Bogenhagen 2001; Takamatsu *et al.* 2002). The core of nucleoid constructed by TFAM-mtDNA, which could also be assembled easily *in vitro* in a droplet-like manner (Farge *et al.* 2014; Feric *et al.* 2021; Long *et al.* 2021). Our data suggest that the nucleoid is self-assembled by phase separation (Long *et al.* 2021), and the flexible linker domain between two HMG domains of TFAM contributes to the phase separation of TFAM *in vitro*. The interaction between TFAM and mtDNA provides another force as the multivalent effect in driving the droplet-like structure of nucleoid. The phase separation promotes the segregation of nucleoids from the mitochondrial matrix, but keeps the interaction between the two of them, which should be a critical feature for mitochondria (Fig. 1).

Phase separation regulates transcription in nuclei

Nuclear transcription should be carried out by multi-component transcription machinery, including hundreds of proteins (Guo *et al.* 2019; Steinbach *et al.* 2019). RNA Polymerase II (Pol II) plays a central role through its polymerase activity in mRNA generation, while the phase separation by the CTD domain also contributes to the recruitment of many other

components for smooth reaction (Boehning *et al.* 2018; Guo *et al.* 2019; Lu *et al.* 2018). This CTD domain contains a conserved repeat of $Y_1S_2P_3T_4S_5P_6S_7$, such as 52 repeats in humans, which served as an IDR for phase separation (Boehning *et al.* 2018). Other subunits necessary for transcription could be further recruited by the co-phase separation manner. The phosphorylation of CTD by CDK7 could switch the transcription from initiation to elongation by regulating the phase separation of Pol II (Boehning *et al.* 2018). The mediators were also reported to act in a similar manner of phase separation and recruit transcription activators to promote transcription or recruit repressors to inhibit transcription activity (Cho *et al.* 2018). The cis-elements, such as enhancers are also reported to be recruited by phase separation, recruiting activators and forming enhancer granules (Sabari *et al.* 2018). Moreover, transcribed RNAs could also regulate the feedback of transcription by phase separation, whereby RNAs initially stimulate condensates of transcription but then ultimately arrest the process by promoting the dissolution of these condensates (Henninger *et al.* 2020). The cleavage of pre-mRNA could also be observed as a droplet-like structure, which may be regulated by phase separation, too (Guo *et al.* 2019) (Fig. 2A).

Phase separation in the modulation of mitochondria transcription

Mitochondria have a series of special machinery in DNA replication and transcription, different from nuclear





Fig. 1 Phase separation in chromatin assembly of the nuclei and the assembly of mitochondrial nucleoid. The schematic diagram represents the phase separation in chromatin condensation and nucleus architecture. The phase separation of TFAM (red) and DNA (gray) drives the nucleoid assembly in mitochondria



The elongation factor TEFM plays critical a role in the transformation of transcription initiation to elongation,

which could also be recruited by co-phase separation with TFAM-DNA. It further remodels the multi-phase separation of POLRMT and promotes the transcription machinery to go further alongside the mtDNA.

The transcription termination in mitochondrial is still not fully determined (Terzioğlu *et al.* 2013; Yakubovskaya *et al.* 2010). The terminal factors as MTERF1 are reported to bind the leu-tRNA site and terminate the transcription from the HSP promoter (Yakubovskaya *et al.* 2010). The MTERF1 could also be recruited by TFAM-DNA in a multi-phase separation manner (Long *et al.* 2021). MTERF1 circling the core complex of the nucleoid is restricted in the outer layer without the touch of most of the mtDNA-TFAM core. This kind of phase separation may also restrict the activity of MTERF1 as non-engagement with the transcription complex. Moreover, MTERF1 could co-phase separation with POLRMT in the outer layer, which may further prevent the activity of free POLRMT in the outer layer.

Similar to the transcription center in nuclei, there are also some special RNA granules in the mitochondria. Mitochondrial RNA granules are clusters of RNA transcripts and ribosome assembling (Pearce *et al.* 2017). The granules known as MRGs, are enrichment with nascent RNAs and RNA-binding proteins (Jourdain *et al.* 2013). It is reported to be an RNA processing center, in which newly synthesized RNAs are cleaved and modified. Many other proteins are also identified in the RNA granules, including GRSF1, FASTKD protein family and so on (Jourdain *et al.* 2013), which may also be driven by phase separation. Moreover, the mitochondrial ribosome assembling process is also a mystery, which may also be happened in a center driven by multi-phase separation, similar to the nucleoids.

SUMMARY AND PERSPECTIVE

The compartmentation by phase separation provides new insight into the architecture and transcription modulation both in nuclei and the mitochondrial nucleoids. Phase separation drives chromatin condensation, nucleolus assembly and nucleus architecture construction. It also drives the mitochondria nucleoid assembly and transcription modulation.

However, there are still many questions that need to be solved. The question of how phase separation regulates chromatin dynamics in cell division and what has happened about the phase separation in control of heterochromatin building and rebuilding process in development is still uncovered. For phase separation in

mitochondria, how does the transcription transform to translation in the phase separation model? Does mtDNA replication share a similar phase separation with transcription? There is still a lot of work to do as a new model and theory, and a lot of questions should be answered.

Acknowledgements This study is funded by the National Natural Science Foundation projects of China (92157202, 32025010, 32241002, 92254301, 32261160376, 31970709, 32070729, 32100619, 32170747), the National Key Research and Development Program of China (2022YFA1103800, 2019YFA0904500), the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB0480000), the Key Research Program, CAS (ZDBS-ZRKJZ-TLC003), International Cooperation Program, CAS (154144KYSB20200006), CAS Project for Young Scientists in Basic Research (YSBR-075), Guangdong Province Science and Technology Program (2023A1515030231, 2022A1515110493, 2020B1212060052, 2021A1515012513, 2021B1515020096, 2022A1515012616, 2022A1515110951), Guangzhou Science and Technology Program (202102020827, 202102080066), Open research funds of the Sixth Affiliated Hospital of Guangzhou Medical University, Qingyuan People's Hospital (202301-203), Open Research Program of Key Laboratory of Regenerative Biology, CAS (KLRB202107, KLRB202203), and CAS Youth Innovation Promotion Association (to Y. W and K. C).

Author contribution X. L. guided the project and wrote the manuscript. Q. L. and Y. Z. wrote the manuscript and prepared the graphs. J. G., H. W. did the manuscript revision.

Compliance with Ethical Standards

Conflict of interest Qi Long, Yanshuang Zhou, Jingyi Guo, Hao Wu and Xingguo Liu declare that they have no conflict of interest.

Human and animal rights and informed consent This article does not contain any studies with human or animal subjects performed by any of the authors.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

Alberti S, Gladfelter A, Mittag T (2019) Considerations and challenges in studying liquid-liquid phase separation and

- biomolecular condensates. *Cell* 176(3): 419–434
- Banani SF, Lee HO, Hyman AA, Rosen MK (2017) Biomolecular condensates: organizers of cellular biochemistry. *Nat Rev Mol Cell Biol* 18(5): 285–298
- Banani SF, Rice AM, Peeples WB, Lin Y, Jain S, Parker R, Rosen MK (2016) Compositional control of phase-separated cellular bodies. *Cell* 166(3): 651–663
- Bi X, Xu Y, Li T, Li X, Li W, Shao W, Wang K, Zhan G, Wu Z, Liu W, Lu JY, Wang L, Zhao J, Wu J, Na J, Li G, Li P, Shen X (2019) RNA targets ribogenesis factor WDR43 to chromatin for transcription and pluripotency control. *Mol Cell* 75(1): 102–116
- Boehning M, Dugast-Darzacq C, Rankovic M, Hansen AS, Yu T, Marie-Nelly H, McSwiggen DT, Kokic G, Dailey GM, Cramer P, Darzacq X, Zweckstetter M (2018) RNA polymerase II clustering through carboxy-terminal domain phase separation. *Nat Struct Mol Biol* 25(9): 833–840
- Boeynaems S, Alberti S, Fawzi NL, Mittag T, Polymenidou M, Rousseau F, Schymkowitz J, Shorter J, Wolozin B, Van Den Bosch L, Tompa P, Fuxreiter M (2018) Protein phase separation: a new phase in cell biology. *Trends Cell Biol* 28(6): 420–435
- Brangwynne CP, Mitchison TJ, Hyman AA (2011) Active liquid-like behavior of nucleoli determines their size and shape in *Xenopus laevis* oocytes. *Proc Natl Acad Sci USA* 108(11): 4334–4339
- Cheng Y, Xie W, Pickering BF, Chu KL, Savino AM, Yang X, Luo H, Nguyen DT, Mo S, Barin E, Velleca A, Rohwetter TM, Patel DJ, Jaffrey SR, Kharas MG (2021) N(6)-Methyladenosine on mRNA facilitates a phase-separated nuclear body that suppresses myeloid leukemic differentiation. *Cancer Cell* 39(7): 958–972
- Cho WK, Spille JH, Hecht M, Lee C, Li C, Grube V, Cisse, II (2018) Mediator and RNA polymerase II clusters associate in transcription-dependent condensates. *Science* 361(6400): 412–415
- Cogliati S, Frezza C, Soriano ME, Varanita T, Quintana-Cabrera R, Corrado M, Cipolat S, Costa V, Casarin A, Gomes LC, Perales-Clemente E, Salviati L, Fernandez-Silva P, Enriquez JA, Scorrano L (2013) Mitochondrial cristae shape determines respiratory chain supercomplexes assembly and respiratory efficiency. *Cell* 155(1): 160–171
- Durniak KJ, Bailey S, Steitz TA (2008) The structure of a transcribing T7 RNA polymerase in transition from initiation to elongation. *Science* 322(5901): 553–557
- Farge G, Mehmedovic M, Baclayon M, van den Wildenberg SM, Roos WH, Gustafsson CM, Wuite GJ, Falkenberg M (2014) *In vitro*-reconstituted nucleoids can block mitochondrial DNA replication and transcription. *Cell Rep* 8(1): 66–74
- Feric M, Demarest TG, Tian J, Croteau DL, Bohr VA, Misteli T (2021) Self-assembly of multi-component mitochondrial nucleoids via phase separation. *EMBO J* 40(6): e107165. <https://doi.org/10.15252/embj.2020107165>
- Feric M, Vaidya N, Harmon TS, Mitrea DM, Zhu L, Richardson TM, Kriwacki RW, Pappu RV, Brangwynne CP (2016) Coexisting liquid phases underlie nucleolar subcompartments. *Cell* 165(7): 1686–1697
- Frey TG, Mannella CA (2000) The internal structure of mitochondria. *Trends Biochem Sci* 25(7): 319–324
- Guo YE, Manteiga JC, Henninger JE, Sabari BR, Dall'Agnese A, Hannett NM, Spille JH, Afeyan LK, Zamudio AV, Shrinivas K, Abraham BJ, Boija A, Decker TM, Rimel JK, Fant CB, Lee TI, Cisse, II, Sharp PA, Taatjes DJ, Young RA (2019) Pol II phosphorylation regulates a switch between transcriptional and splicing condensates. *Nature* 572(7770): 543–548
- Gustafsson CM, Falkenberg M, Larsson NG (2016) Maintenance and expression of mammalian mitochondrial DNA. *Annu Rev Biochem* 85: 133–160
- Hansen AS, Hsieh TS, Cattoglio C, Pustova I, Saldana-Meyer R, Reinberg D, Darzacq X, Tjian R (2019) Distinct classes of chromatin loops revealed by deletion of an RNA-binding region in CTCF. *Mol Cell* 76(3): 395–411
- Henninger JE, Oksuz O, Shrinivas K, Sagi I, LeRoy G, Zheng MM, Andrews JO, Zamudio AV, Lazaris C, Hannett NM, Lee TI, Sharp PA, Cisse, II, Chakraborty AK, Young RA (2020) RNA-mediated feedback control of transcriptional condensates. *Cell* 184(1): 207–225
- Hillen HS, Morozov YI, Sarfallah A, Temiakov D, Cramer P (2017) Structural basis of mitochondrial transcription initiation. *Cell* 171(5): 1072–1081
- Hillen HS, Temiakov D, Cramer P (2018) Structural basis of mitochondrial transcription. *Nat Struct Mol Biol* 25(9): 754–765
- Jourdain AA, Koppen M, Wydro M, Rodley CD, Lightowlers RN, Chrzanowska-Lightowlers ZM, Martinou JC (2013) GRSF1 regulates RNA processing in mitochondrial RNA granules. *Cell Metab* 17(3): 399–410
- Klein IA, Boija A, Afeyan LK, Hawken SW, Fan M, Dall'Agnese A, Oksuz O, Henninger JE, Shrinivas K, Sabari BR, Sagi I, Clark VE, Platt JM, Kar M, McCall PM, Zamudio AV, Manteiga JC, Coffey EL, Li CH, Hannett NM, Guo YE, Decker TM, Lee TI, Zhang T, Weng JK, Taatjes DJ, Chakraborty A, Sharp PA, Chang YT, Hyman AA, Gray NS, Young RA (2020) Partitioning of cancer therapeutics in nuclear condensates. *Science* 368(6497): 1386–1392
- Kukat C, Davies KM, Wurm CA, Spähr H, Bonekamp NA, Köhl I, Joos F, Polosa PL, Park CB, Posse V, Falkenberg M, Jakobs S, Köhlbrandt W, Larsson NG (2015) Cross-strand binding of TFAM to a single mtDNA molecule forms the mitochondrial nucleoid. *Proc Natl Acad Sci USA* 112(36): 11288–11293
- Ladouceur AM, Parmar BS, Biedzinski S, Wall J, Tope SG, Cohn D, Kim A, Soubry N, Reyes-Lamothe R, Weber SC (2020) Clusters of bacterial RNA polymerase are biomolecular condensates that assemble through liquid-liquid phase separation. *Proc Natl Acad Sci USA* 117(31): 18540–18549
- Lafontaine DLJ, Riback JA, Bascetin R, Brangwynne CP (2021) The nucleolus as a multiphase liquid condensate. *Nat Rev Mol Cell Biol* 22(3): 165–182
- Larson AG, Elnatan D, Keenen MM, Trnka MJ, Johnston JB, Burlingame AL, Agard DA, Redding S, Narlikar GJ (2017) Liquid droplet formation by HP1 α suggests a role for phase separation in heterochromatin. *Nature* 547(7662): 236–240
- Larson AG, Narlikar GJ (2018) The role of phase separation in heterochromatin formation, function, and regulation. *Biochemistry* 57(17): 2540–2548
- Lee JH, Wang R, Xiong F, Krakowiak J, Liao Z, Nguyen PT, Moroz-Omori EV, Shao J, Zhu X, Bolt MJ, Wu H, Singh PK, Bi M, Shi CJ, Jamal N, Li G, Mistry R, Jung SY, Tsai KL, Ferreon JC, Stossi F, Cafilisch A, Liu Z, Mancini MA, Li W (2021) Enhancer RNA m6A methylation facilitates transcriptional condensate formation and gene activation. *Mol Cell* 81(16): 3368–3385
- Long Q, Zhou Y, Wu H, Du S, Hu M, Qi J, Li W, Guo J, Wu Y, Yang L, Xiang G, Wang L, Ye S, Wen J, Mao H, Wang J, Zhao H, Chan WY, Liu J, Chen Y, Li P, Liu X (2021) Phase separation drives the self-assembly of mitochondrial nucleoids for transcriptional modulation. *Nat Struct Mol Biol* 28(11): 900–908
- Lu H, Yu D, Hansen AS, Ganguly S, Liu R, Heckert A, Darzacq X, Zhou Q (2018) Phase-separation mechanism for C-terminal hyperphosphorylation of RNA polymerase II. *Nature* 558(7709): 318–323
- Masters BS, Stohl LL, Clayton DA (1987) Yeast mitochondrial RNA

- polymerase is homologous to those encoded by bacteriophages T3 and T7. *Cell* 51(1): 89–99
- Misteli T (2020) The self-organizing genome: principles of genome architecture and function. *Cell* 183(1): 28–45
- Nava MM, Miroshnikova YA, Biggs LC, Whitefield DB, Metge F, Boucas J, Vihinen H, Jokitalo E, Li X, Garcia Arcos JM, Hoffmann B, Merkel R, Niessen CM, Dahl KN, Wickstrom SA (2020) Heterochromatin-driven nuclear softening protects the genome against mechanical stress-induced damage. *Cell* 181(4): 800–817
- Ngo HB, Lovely GA, Phillips R, Chan DC (2014) Distinct structural features of TFAM drive mitochondrial DNA packaging versus transcriptional activation. *Nat Commun* 5: 3077. <https://doi.org/10.1038/ncomms4077>
- Pearce SF, Rebelo-Guiomar P, D'Souza AR, Powell CA, Van Haute L, Minczuk M (2017) Regulation of mammalian mitochondrial gene expression: recent advances. *Trends Biochem Sci* 42(8): 625–639
- Riback JA, Zhu L, Ferrolino MC, Tolbert M, Mitrea DM, Sanders DW, Wei M-T, Kriwacki RW, Brangwynne CP (2020) Composition-dependent thermodynamics of intracellular phase separation. *Nature* 581(7807): 209–214
- Roger AJ, Munoz-Gomez SA, Kamikawa R (2017) The origin and diversification of mitochondria. *Curr Biol* 27(21): R1177–R1192
- Sabari BR, Dall'Agnese A, Boija A, Klein IA, Coffey EL, Shrinivas K, Abraham BJ, Hannett NM, Zamudio AV, Manteiga JC, Li CH, Guo YE, Day DS, Schuijers J, Vasile E, Malik S, Hnisz D, Lee TI, Cisse, II, Roeder RG, Sharp PA, Chakraborty AK, Young RA (2018) Coactivator condensation at super-enhancers links phase separation and gene control. *Science* 361(6400): eaar3958. <https://doi.org/10.1126/science.aar3958>
- Sabari BR, Dall'Agnese A, Young RA (2020) Biomolecular condensates in the nucleus. *Trends Biochem Sci* 45(11): 961–977
- Shen EL, Bogenhagen DF (2001) Developmentally-regulated packaging of mitochondrial DNA by the HMG-box protein mtTFA during *Xenopus oogenesis*. *Nucleic Acids Res* 29(13): 2822–2828
- Shin Y, Brangwynne CP (2017) Liquid phase condensation in cell physiology and disease. *Science* 357(6357): eaaf4382. <https://doi.org/10.1126/science.aaf4382>
- Song F, Chen P, Sun D, Wang M, Dong L, Liang D, Xu RM, Zhu P, Li G (2014) Cryo-EM study of the chromatin fiber reveals a double helix twisted by tetranucleosomal units. *Science* 344(6182): 376–380
- Steinbach N, Hasson D, Mathur D, Stratikopoulos EE, Sachidanandam R, Bernstein E, Parsons RE (2019) PTEN interacts with the transcription machinery on chromatin and regulates RNA polymerase II-mediated transcription. *Nucleic Acids Res* 47(11): 5573–5586
- Strickfaden H, Tolsma TO, Sharma A, Underhill DA, Hansen JC, Hendzel MJ (2020) Condensed chromatin behaves like a solid on the mesoscale *in vitro* and in living cells. *Cell* 183(7): 1772–1784
- Strom AR, Emelyanov AV, Mir M, Fyodorov DV, Darzacq X, Karpen GH (2017) Phase separation drives heterochromatin domain formation. *Nature* 547(7662): 241–245
- Takamatsu C, Umeda S, Ohsato T, Ohno T, Abe Y, Fukuoh A, Shinagawa H, Hamasaki N, Kang D (2002) Regulation of mitochondrial D-loops by transcription factor A and single-stranded DNA-binding protein. *EMBO Rep* 3(5): 451–456
- Terzioglu M, Ruzzenente B, Harmel J, Mourier A, Jemt E, Lopez MD, Kukut C, Stewart JB, Wibom R, Meharg C, Habermann B, Falkenberg M, Gustafsson CM, Park CB, Larsson NG (2013) MTERF1 binds mtDNA to prevent transcriptional interference at the light-strand promoter but is dispensable for rRNA gene transcription regulation. *Cell Metab* 17(4): 618–626
- Wang J, Choi JM, Holehouse AS, Lee HO, Zhang X, Jahnel M, Maharana S, Lemaitre R, Pozniakovsky A, Drechsel D, Poser I, Pappu RV, Alberti S, Hyman AA (2018) A molecular grammar governing the driving forces for phase separation of prion-like RNA binding proteins. *Cell* 174(3): 688–699
- Wang L, Hu M, Zuo MQ, Zhao J, Wu D, Huang L, Wen Y, Li Y, Chen P, Bao X, Dong MQ, Li G, Li P (2020) Rett syndrome-causing mutations compromise MeCP2-mediated liquid-liquid phase separation of chromatin. *Cell Res* 30(5): 393–407
- Yakubovskaya E, Mejia E, Byrnes J, Hambardjiev E, Garcia-Diaz M (2010) Helix unwinding and base flipping enable human MTERF1 to terminate mitochondrial transcription. *Cell* 141(6): 982–993